arXiv:1303.6355v1 [hep-ph] 26 Mar 2013

Decoding the riddle of $Y(4260)$ and $Z_c(3900)$

Qian Wang[1*], Christoph Hanhart[1†], Qiang Zhao[2‡]
[1] *Institut für Kernphysik and Institute for Advanced Simulation,*
Forschungszentrum Jülich, D–52425 Jülich, Germany
[2] *Institute of High Energy Physics and Theoretical Physics Center for Science Facilities,*
Chinese Academy of Sciences, Beijing 100049, China

(Dated: March 27, 2013)

The observation of $Z_c(3900)$ by the BESIII collaboration in the invariant mass spectrum of $J/\psi\pi^\pm$ in $e^+e^- \to J/\psi\pi^+\pi^-$ at the center of mass 4.260 GeV suggests the existence of a charged $\bar{D}D^*+D\bar{D}^*$ molecular state with $I(J^P) = 1(1^+)$, which could be an isovector brother of the famous $X(3872)$ and an analogue of $Z_b(10610)$ claimed by the Belle Collaboration. We demonstrate that this observation provides strong evidence that the mysterious $Y(4260)$ is a $\bar{D}D_1(2420)+D\bar{D}_1(2420)$ molecular state. Especially, we show that the decay of this molecule naturally populates low momentum $\bar{D}D^*$ pairs and leads unavoidably to a cusp at the $\bar{D}D^*$ threshold. We discuss the signatures that distinguish such a $\bar{D}D^*$ cusp from the presence of a true resonance.

PACS numbers: 14.40.Rt, 13.75.Lb, 13.20.Gd

During the past years, the experimental observation of a large number of so-called X, Y, Z states has initiated tremendous efforts to unravel their nature beyond the conventional quark model. Especially, the conformation of signals in charged channels would be a direct evidence for exotic states. For instance, the Belle Collaboration reported signals for $Z(4430)$ in $\psi'\pi^\pm$, and $Z_1(4050)$ and $Z_2(4250)$ in $\chi_{c1}\pi^\pm$ in B meson decays [1]. However, an enhancement in the same mass range was interpreted as a reflection by BaBar [2]. The more recent experimental results for charged bottomonium states $Z_b(10610)$ and $Z_b(10650)$, located close to the $\bar{B}B^*$ and $\bar{B}^*B^*$ thresholds, respectively, by the Belle Collaboration [3] seem to be the first strong evidence for QCD "exotics" in the heavy quark sector. In this context the recent report of an enhancement in the $J/\psi\pi^\pm$ invariant mass distribution around 3900 MeV, right at the $\bar{D}D^*$ threshold, by the BESIII collaboration [4] clearly reinforces the existence of such an unusual phenomenon. This state, called $Z_c(3900)$ below, might be an isovector partner of the well established 1^{++} isoscalar $X(3872)$ [5], but with $I^G(J^{PC}) = 1^+(1^{+-})$ for the neutral state.

In this work we demonstrate that, if $Y(4260)$ is a $\bar{D}D_1 + c.c.$ molecule (below we use $\bar{D}D_1$ as a short notation), the appearance of an enhancement around 3900 MeV in the $J/\psi\pi$ invariant mass distribution can be shown to be natural. Here D_1 refers to the narrow axial vector $D_1(2420)$ ($\Gamma = 27 \pm 3$ MeV) with $I(J^P) = \frac{1}{2}(1^+)$ [5]. In this sense the observation of the charged $Z_c(3900)$ state by BESIII in $Y(4260) \to J/\psi\pi\pi$ provides a very strong evidence for the molecular nature of $Y(4260)$. We also discuss whether the observed enhancement can be interpreted purely as a cusp or whether the inclusion of explicit poles in the $\bar{D}D^*$ system is necessary. Before we go into details of our calculations we first briefly review the status of $Y(4260)$.

The $Y(4260)$ was observed via its decays into $J/\psi\pi^+\pi^-$, $J/\psi\pi^0\pi^0$, and $J/\psi K^+K^-$ by various experiments. Its mass and total width are well determined as $m = 4263^{+8}_{-9}$ MeV and $\Gamma_Y = 95 \pm 14$ MeV, respectively [5]. The new data from BESIII confirms the signal in $Y(4260) \to J/\psi\pi^+\pi^-$ with much higher statistics. The most mysterious fact about $Y(4260)$ is not that its mass does not agree to what is predicted by the potential quark model. Instead, as a charmonium state with $J^{PC} = 1^{--}$, it is only "seen" as a bump in its two pion transitions to J/ψ, but not in any open charm decay channel like $\bar{D}D$, $\bar{D}D^* + c.c.$, $\bar{D}^*D^*$ and $\bar{D}_sD_s^*$, or other tens of measured channels. In fact, the cross section lineshapes of the e^+e^- annihilations into $D^{(*)}$ meson pairs appear to have a dip at its peak mass 4.26 GeV instead of a bump.

In the vector sector with $J^{PC} = 1^{--}$, one should recognize that the $\bar{D}D_1(2420)$ is the first open charm relative S-wave channel that is coupled to this quantum number, and the nominal threshold is only about 29 MeV above the location of the $Y(4260)$. The possibility that $Y(4260)$ may be a bound system of $\bar{D}D_1$ and $\bar{D}^*D_0$ was investigated in Refs. [6–9]. It should be stressed, however, that broad components (the widths of D_0 and $D_1(2430)$ are as large as 300 MeV) can not produce narrow resonances [10]. We thus do not consider the $\bar{D}^*D_0$ or $\bar{D}^*D_1(2430)$ component here, but focus on the assumption that the $Y(4260)$ is a bound system of only $\bar{D}D_1(2420)$.

---

[*] *Email address:* q.wang@fz-juelich.de
[†] *Email address:* c.hanhart@fz-juelich.de
[‡] *Email address:* zhaoq@ihep.ac.cn



FIG. 1: The Feynman diagrams for $Y(4260) \to \bar{D}D_1 + c.c. \to J/\psi\pi\pi$ and $h_c\pi\pi$ considered in this work.

It should be emphasized that in the literature many other solutions were proposed for the $Y(4260)$ (see Ref. [11] for a recent review). From a fit to the $\pi\pi$ spectrum of $Y(4260) \to J/\psi\pi\pi$ from the BaBar Collaboration [12], Dai *et al.* [13] concluded that the even lower threshold $\chi_{c0}\omega$ would have the largest coupling to $Y(4260)$, while the $\bar{D}D_1$ coupling to $Y(4260)$ turned out to be negligible. However, the observation of the enhancement at the $\bar{D}D^*$ threshold in the $J/\psi\pi$ invariant mass spectrum actually rules out such a scenario and suggests that the underlying dynamics should be more sensitive to the $\bar{D}D_1$ threshold.

In this work we do not try to identify the mechanisms that lead to the formation of the $Y(4260)$ as a molecular state, but study the consequences of the assumption that this state is a $\bar{D}D_1$ bound system. We argue that the interpretation of $Y(4260)$ as a relative S-wave $\bar{D}D_1$ system is able to accommodate nearly all the present observations for $Y(4260)$. Especially, its absence in various open charm decay channels mentioned above and the observation of $Z_c(3900)$ in $Y(4260) \to J/\psi\pi\pi$ can be naturally understood.

The heavy quark spin symmetry also implies the presence of a $\bar{D}^*D_1$ and $\bar{D}^*D_2$ component in the wave function. However, they will be neglected here since their corresponding thresholds are almost 200 MeV above the mass of $Y(4260)$. Meanwhile, we note that in a more microscopic treatment there should be heavier spin partners of $Y(4260)$ that should contain the mentioned constituents prominently. We will briefly come back to this issue at the end of this paper.

We now provide the details of our calculations based on the picture described above. The pertinent diagrams are shown in Fig. 1. The central goal of our study is to pin down the structure of $Y(4260)$ and identify the quantitative importance of the $\bar{D}D^*$ cusp from the loop diagrams (b)-(d) in order to understand whether the BESIII spectra call for the additional inclusion of an explicit pole diagram as depicted in diagram (a).

It should be stressed that the vicinity of the $\bar{D}D_1$ threshold to 4.26 GeV favors the formation of low momentum $\bar{D}D^*$ pairs, which may lead to the formation of the $Z_c(3900)$ bound systems. The reason is that the $\bar{D}D_1$ intermediate state as well as the $\bar{D}D^*$ intermediate state can be simultaneously close to their mass shells — all with relative S waves. This gives rise to a triangle singularity studied in a different context in Refs. [14, 15]. Such a two-cut condition strongly enhances the corresponding matrix elements. In this sense the $\bar{D}D_1$ intermediate system provides an ideal doorway state for a low momentum $\bar{D}D^*$ system. There are a series of S-wave open charm thresholds with $J^{PC} = 1^{--}$ around 4.26 GeV, i.e. $\bar{D}D_1(2420)$, $\bar{D}D_1(2430)$, $\bar{D}^*D_0$, $\bar{D}_{s0}D_s^*$, $\bar{D}_sD_{s1}$, and $\bar{D}^*D_1(2420)$. However, all of them, except for $\bar{D}D_1(2420)$, are either far away from the observed physical $Y(4260)$, or too broad to make a bound state [5]. Thus, we do not take them into account explicitly here.

Note that in order to formulate the problem as an effective field theory the power counting of Refs. [16, 17] needs to be adapted to the present situation. We leave this to be reported in a subsequent work and focus here on a more phenomenological investigation.

A complete calculation would also need the inclusion of the $\pi\pi$ final state interaction. This should be done in a way consistent with the Watson theorem. Thus, an inclusion of final state interactions in all diagrams is needed. In addition, since the pion pairs are in the isoscalar channel, also $\bar{K}K$ intermediate states need to be considered. This is beyond the scope of this paper and will be studied in a followed-up work. However, as a result of the omission of the $\pi\pi$ final state interaction we do not expect to be able to properly describe the $\pi\pi$ invariant mass distribution observed by BESIII.



FIG. 2: The invariant mass spectra for (a) $J/\psi\pi$ and (b) $\pi\pi$ in $Y(4260) \to J/\psi\pi\pi$. The solid, dashed, and dot-dot-dashed lines corresponds to the results of the full calculation, box diagrams, and triangle diagram with the $Z_c(3900)$ pole, respectively.

Since $Z_c(3900)$ has the same quantum numbers as Z_b most interactions needed for this work can be taken from Ref. [17]. Here we only present the interactions between a P-wave charmed meson and other fields since they play a crucial role in this work. The heavy quark spin symmetry allows heavy mesons to form spin doublet super fields distinguished by their light degrees of freedom $s_l = s_q + l$ with s_q the light quark spin and l the orbital angular momentum. For $l = 1$ it can be classified into two super fields [18, 19]. The interaction terms relevant for this work read

$$\begin{aligned}
\mathcal{L}_Y &= iy(\bar{D}_a^\dagger Y^i D_{1a}^{i\dagger} - \bar{D}_{1a}^{i\dagger} Y^i D_a^\dagger) + H.c., &(1)\\
\mathcal{L}_{D_1} &= i\frac{h'}{f_\pi}[6D_{1a}^i(\partial^i\partial^j\phi_{ab})D_b^{*\dagger j} - D_{1a}^i(\partial^j\partial^j\phi_{ab})D_b^{*\dagger i} \\
&\quad -6\bar{D}_a^{*\dagger i}(\partial^i\partial^j\phi_{ab})\bar{D}_{1b}^j + \bar{D}_a^{*\dagger i}(\partial^j\partial^j\phi_{ab})\bar{D}_{1b}^i] + H.c., &(2)
\end{aligned}$$

where D ($D^\dagger$) and $\bar{D}$ ($\bar{D}^\dagger$) contain the annihilation (creation) operators for $c\bar{q}$ and $\bar{c}q$ fields, respectively. The analogous conventions are applied to D^* and D_1. The interactions contain five coupling constants in total. Since we focus on the shape of the invariant mass distributions only, all couplings that are common to all diagrams are not relevant. In this sense the only free parameters that influence the invariant mass distributions are the mass of $Z_c(3900)$ and the coupling constant z for $Z_c\bar{D}D^*$.

In analogy to what is known about the Z_b states we assume that the width of the $Z_c(3900)$ is saturated by the $\bar{D}D^*$ intermediate state. Then the width of Z_c that enters the Flatte distribution is controlled by the same parameter z introduced above.

For simplicity in this exploratory study we treat $Y(4260)$, $D_1(2420)$, D and D^* as stable states. Their masses are taken from the Particle Data Group [5]. For the $Z_c(3900)$ propagator we use a Flatte distribution, as mentioned above. By assuming that $Y(4260)$ is dominated by the $\bar{D}D_1(2420)$ molecule component, we can estimate the YDD_1 coupling by Weinberg's compositeness theorem [20, 21], i.e. $y^2/4\pi \equiv 4(m_D + m_{D_1})^{5/2}\sqrt{2\delta E/m_D m_{D_1}} \simeq 17\ \text{GeV}^2$. This predicts the dominant decay of $Y(4260) \to \bar{D}D^*\pi + c.c.$ via the intermediate state $\bar{D}D_1(2420) + c.c.$ of which the partial width is larger than 40 MeV. This value is consistent with the total width measured for $Y(4260)$. In addition, the $\bar{D}D_1$ component should also distort the invariant mass distributions for $Y(4260) \to \bar{D}D^*\pi + c.c.$ A more quantitative result will be reported in a subsequent work.

The numerical results for the $J/\psi\pi$ and $\pi\pi$ invariant mass spectra from $Y(4260) \to J/\psi\pi\pi$ are shown in Fig. 2. The results from the box diagrams (Fig. 1 (b)-(d)) are shown as the dashed lines, while the result for the $Z_c(3900)$ pole diagrams, Fig. 1 (a), is shown as the dot-dot-dashed lines. The full result, from the interference of these two contributions, is shown as the solid lines. One can see that even without the explicit inclusion of $Z_c(3900)$, two structures appear at the same masses in the $J/\psi\pi$ spectrum as in the BESIII data. Note that by charge conjugation invariance the $J/\psi\pi^-$ spectrum is identical with that for $J/\psi\pi^+$. The cusp at $M_D + M_{D^*} \simeq 3.876$ GeV in Fig. 2(a) marks the $\bar{D}D^*$ threshold. The lower bump between 3.4 and 3.6 GeV in the $J/\psi\pi^+$ invariant mass distribution comes from the interference between the conjugate diagrams where either a π^- or a π^+ is emitted first. Therefore, the lower bump is simply a reflection of the narrow structure at 3.9 GeV.

Although one also observes that the agreement of the dashed curve in Fig. 2 with the data is only qualitative, the appearance of a visible cusp supports the claim made in the introduction that the loop kinematics favors the production of low momentum $\bar{D}D^*$ pairs. However, the effect of the cusp alone is too small and too narrow compared to the data. This implies that the experimental observation of a very pronounced, broader peak at the $\bar{D}D^*$ threshold suggests the need for an explicit resonance pole on top of the cusp. In this sense, the BESIII results seem to have confirmed the existence of the isovector partner of $X(3872)$.



FIG. 3: The invariant mass spectra for (a) $h_c\pi$ and (b) $\pi\pi$ in $Y(4260) \to h_c\pi\pi$. The notations are the same as in Fig. 2.

The solid lines in Fig. 2 show the result of the sum of all diagrams in Fig. 1. The Flatte parameters are chosen in order to reproduce the qualitative features of the data. We find that with $M_{Z_c} = 3.92$ GeV and $z = 12$ GeV$^{-\frac{1}{2}}$ the main features of the data can be reproduced. We postpone a more refined extraction of the parameters, including the uncertainties, until the $\pi\pi$ final state interaction is included properly.

As expected, the $\pi\pi$ spectrum of Fig. 2 (b) is in disagreement with the data, however, we expect that this discrepancy can be overcome, once we include the $\pi\pi$ final state interaction.

In Fig. 1, we also included the production channel for $Y(4260) \to h_c\pi\pi$. Similar to the discussions of $Z_b \to h_b\pi$, this channel is ideal for disentangling the molecular nature of the intermediate $Z_c(3900)$: The power counting analysis in Ref. [17] shows that the triangle transition $Z_b \to h_b(mP)\pi$ is not suppressed compared to $Z_b \to \Upsilon(nS)\pi$, although the decay is via a P-wave. This explains why the branching ratios for $Z_b \to h_b(mP)\pi$ are compatible with those for $Z_b \to \Upsilon(nS)\pi$. A similar phenomenon occurs here. In addition, higher loop contributions are suppressed in $Z_b \to h_b(mP)\pi$, while they are not suppressed for $Z_b \to \Upsilon\pi$ [17]. The analogous pattern is expected for the Z_c decays. However, as we will see, the box diagrams contribute negligibly to the transition $Z_c \to h_c\pi$ such that we are still able to predict the shape of the $h_c\pi$ invariant mass distributions. Therefore, a measurement of $Z_c \to h_c\pi$ and its comparison with $Z_c \to J/\psi\pi$ will provide a further non-trivial confirmation of the molecular nature of $Z_c(3900)$.

In Fig. 3, we present our prediction for the invariant mass spectra of $h_c\pi$ and $\pi\pi$ in $Y(4260) \to h_c\pi\pi$. The notation of the curves is the same as in Fig. 2. Similar to the $J/\psi\pi\pi$ channel, a very pronounced peak right at the $\bar{D}D^*$ threshold appears in the $h_c\pi$ invariant spectrum. Interestingly, due to the limited phase space in this decay channel, its kinematic reflection is significantly shifted and located at higher invariant masses. As mentioned, for the $h_c\pi$ final state the contribution from the box diagram appears to be negligible. Thus, once measured, this channel will allow a clean experimental access to the $Z_c(3900)$ resonance contribution.

We stress that in order to understand the gross features of the data the diagrams in Fig. 1 can also be replaced completely or in parts by tree level diagrams (see, e.g., Refs. [22, 23]). Then, however, the underlying dynamics with the interplay of the $\bar{D}D^*$ and $\bar{D}D_1$ cuts will be lost. We reiterate that it is the $\bar{D}D_1$ molecule nature of $Y(4260)$ that provides a natural explanation for the appearance of $Z_c(3900)$ in the $Y(4260)$ decays, and for other detailed features of the spectra.

In this work we propose that $Y(4260)$ is dominantly a $\bar{D}D_1(2420)$ molecule and identify a unique mechanism, namely the presence of two-cut condition, which plays an essential role in $Y(4260) \to J/\psi\pi\pi$ and $h_c\pi\pi$. We demonstrate that without introducing any drastic assumption, the molecular nature of $Y(4260)$ as a bound state of $\bar{D}D_1$ can naturally explain the observation of an enhancement around 3.876 GeV in the $J/\psi\pi$ invariant mass spectrum. The reflection of this peak in the invariant mass spectrum produces a lower mass bump around 3.5 GeV. This signature matches the experimental data, thus, provides strong evidence for the molecular nature of $Y(4260)$. We also demonstrate that for a more detailed description of the data it appears necessary to include explicit $Z_c(3900)$ poles, i.e. a resonance structure as an isovector partner of $X(3872)$.

We stress the following important consequences of this prescription that will be reported in a subsequent paper:

i) Although the nominal $\bar{D}D_1(2420)$ threshold is higher than the mass of $Y(4260)$, the threshold of $\bar{D}D^*\pi$ (then decaying to $\bar{D}D\pi\pi$) is much lower. Dominant decays of $Y(4260)$ into $\bar{D}D^*\pi$ should be regarded as a natural consequence and would explain the large deficit in width between the total width and its decays into $J/\psi\pi\pi$ and $J/\psi K\bar{K}$. We also predict an asymmetric spectral shape for the decay of $Y(4260)$ into $\bar{D}D^*\pi$. This can be clarified by an energy scan around the nominal Y mass (analogous to what happens to the shape of $Y(4660)$ when being viewed as $\psi' f_0(980)$ molecule [24]).

ii) A possible reason why the isovector partner $Z_c(3900)$ of the $X(3872)$ was not observed in B decays is that the $X(3872)$ might be produced via its small $\bar{c}c$ component which is absent in Z_c. Here, $Z_c(3900)$ does show up as a result of the proposed molecular nature of $Y(4260)$, which actually allows the S-wave $\bar{D}D^*$ pairs to be copiously produced.

iii) One might also expect a cusp or possible resonance structure at the $\bar{D}^*D^*$ threshold in the decay of $\psi(4415)$, if this state is assumed to be a molecule driven by the nearby $\bar{D}^*D_1$ threshold.

iv) The exact mechanism at work here might also be the reason for the appearance of the Z_b and Z_b' states in $\Upsilon(5S)$ decays, if we assume that $\Upsilon(5S)$ has a sizeable $\bar{B}B_1$ component. Note that the relative S-wave threshold for $\bar{B}B_1$ is only about 120 MeV above the $\Upsilon(5S)$ mass.

The authors thank F.-K. Guo, G. Li, U.-G. Meissner, C.-Z. Yuan, and B.-S. Zou for useful discussions. This work is supported, in part, by the National Natural Science Foundation of China (Grant Nos. 11035006 and 11121092), the Chinese Academy of Sciences (KJCX3-SYW-N2), the Ministry of Science and Technology of China (2009CB825200), and DFG and NSFC funds to the Sino-German CRC 110.

---

[1] R. Mizuk *et al.* [Belle Collaboration], Phys. Rev. D **78** (2008) 072004.
[2] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **79** (2009) 112001.
[3] A. Bondar *et al.* [Belle Collaboration], Phys. Rev. Lett. **108** (2012) 122001.
[4] M. Ablikim *et al.*, [BESIII Collaboration], arXiv:1303.5949 [hep-ex].
[5] J. Beringer *et al.* [Particle Data Group Collaboration], Phys. Rev. D **86**, 010001 (2012).
[6] E. Swanson, Int. J. Mod. Phys. A **21** (2006) 733 [AIP Conf. Proc. **814** (2006) 203].
[7] F.E. Close *et al.*, Phys. Rev. D **81**, 074033 (2010).
[8] G. -J. Ding, Phys. Rev. D **79**, 014001 (2009).
[9] M. -T. Li, W. -L. Wang, Y. -B. Dong and Z. -Y. Zhang, arXiv:1303.4140 [nucl-th].
[10] A. A. Filin *et al.*, Phys. Rev. Lett. **105** (2010) 019101.
[11] N. Brambilla, *et al.*, Eur. Phys. J. C **71**, 1534 (2011).
[12] J. P. Lees *et al.* [BABAR Collaboration], Phys. Rev. D **86**, 051102 (2012).
[13] L. Y. Dai *et al.*, arXiv:1206.6911 [hep-ph].
[14] J. -J. Wu *et al.*, Phys. Rev. Lett. **108**, 081803 (2012).
[15] X. -G. Wu *et al.* Phys. Rev. D **87**, 014023 (2013).
[16] F. -K. Guo *et al.* , Phys. Rev. D **83**, 034013 (2011).
[17] M. Cleven *et al.*, arXiv:1301.6461 [hep-ph], Phys. Rev. D in press.
[18] R. Casalbuoni *et al.* , Phys. Rept. **281**, 145 (1997).
[19] P. Colangelo *et al.* , Phys. Lett. B **634**, 235 (2006).
[20] S. Weinberg, Phys. Rev. **137** (1965) B672.
[21] V. Baru *et al.*, Phys. Lett. B **586**, 53 (2004).
[22] D. -Y. Chen, X. Liu and T. Matsuki, arXiv:1208.2411 [hep-ph].
[23] D. -Y. Chen and X. Liu, Phys. Rev. D **84**, 094003 (2011).
[24] F. -K. Guo *et al.*, Phys. Lett. B **665**, 26 (2008).